

# EPLERENONE- eplerenone tablet, film coated

## Greenstone LLC

### HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use eplerenone tablets safely and effectively. See full prescribing information for eplerenone tablets.

#### EPLERENONE tablets

Initial U.S. Approval: 2002

### INDICATIONS AND USAGE

Eplerenone is an aldosterone antagonist indicated for:

- Improving survival of stable patients with LV systolic dysfunction (LVEF  $\leq 40\%$ ) and CHF after an acute myocardial infarction. (1.2)
- Hypertension, alone or combined with other agents. (1.3)

Eplerenone is indicated for the treatment of hypertension, to lower blood pressure. Lowering blood pressure reduces the risk of fatal and nonfatal cardiovascular events, primarily strokes and myocardial infarctions.

### DOSAGE AND ADMINISTRATION

**CHF Post-MI:** Initiate treatment with 25 mg once daily. Titrate to maximum of 50 mg once daily within 4 weeks, as tolerated. Dose adjustments may be required based on potassium levels. (2.1)

**Hypertension:** 50 mg once daily, alone or combined with other antihypertensive agents. For inadequate response, increase to 50 mg twice daily. Higher dosages are not recommended. (2.2)

**For all patients:**

Measure serum potassium before starting eplerenone and periodically thereafter. (2.3)

### DOSAGE FORMS AND STRENGTHS

Tablets: 25 mg, 50 mg (3)

### CONTRAINDICATIONS

**For all patients:**

- Serum potassium  $> 5.5$  mEq/L at initiation (4)
- Creatinine clearance  $\leq 30$  mL/min (4)
- Concomitant use with strong CYP3A4 inhibitors (4, 7.1)

**For the treatment of hypertension:**

- Type 2 diabetes with microalbuminuria (4)
- Serum creatinine  $> 2.0$  mg/dL in males,  $> 1.8$  mg/dL in females (4)
- Creatinine clearance  $< 50$  mL/min (4)
- Concomitant use of potassium supplements or potassium-sparing diuretics (4)

### WARNINGS AND PRECAUTIONS

- Hyperkalemia: Patients with decreased renal function and diabetics with proteinuria are at increased risk. Proper patient selection and monitoring and avoiding certain concomitant medications can minimize the risk. (5.1)

### ADVERSE REACTIONS

**CHF Post-MI:** Most common adverse reactions ( $> 2\%$  and more frequent than with placebo): hyperkalemia and increased creatinine. (6.1)

**Hypertension:** Most common adverse reactions ( $\geq 2\%$  and more frequent than with placebo): dizziness, diarrhea, coughing, fatigue and flu-like symptoms. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Greenstone LLC at 1-800-438-1985 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).

### DRUG INTERACTIONS

- CYP3A4 Inhibitors: Reduce the starting dose for hypertension to 25 mg once daily when used with moderate CYP3A4 inhibitors (e.g., verapamil, erythromycin, saquinavir, fluconazole). (2.4, 7.1, 12.3)

See 17 for PATIENT COUNSELING INFORMATION.

Revised: 11/2014

---

## **FULL PRESCRIBING INFORMATION: CONTENTS\***

### **1 INDICATIONS AND USAGE**

- 1.1 Patient Selection Considerations
- 1.2 Congestive Heart Failure Post-Myocardial Infarction
- 1.3 Hypertension

### **2 DOSAGE AND ADMINISTRATION**

- 2.1 Congestive Heart Failure Post-Myocardial Infarction
- 2.2 Hypertension
- 2.3 Recommended Monitoring
- 2.4 Dose Modifications for Specific Populations

### **3 DOSAGE FORMS AND STRENGTHS**

### **4 CONTRAINDICATIONS**

### **5 WARNINGS AND PRECAUTIONS**

- 5.1 Hyperkalemia
- 5.2 Impaired Hepatic Function
- 5.3 Impaired Renal Function

### **6 ADVERSE REACTIONS**

- 6.1 Clinical Trials Experience
- 6.2 Clinical Laboratory Test Findings
- 6.3 Postmarketing Experience

### **7 DRUG INTERACTIONS**

- 7.1 CYP3A4 Inhibitors
- 7.2 ACE Inhibitors and Angiotensin II Receptor Antagonists
- 7.3 Lithium
- 7.4 Nonsteroidal Anti-Inflammatory Drugs (NSAIDs)

### **8 USE IN SPECIFIC POPULATIONS**

- 8.1 Pregnancy
- 8.3 Nursing Mothers
- 8.4 Pediatric Use
- 8.5 Geriatric Use

### **10 OVERDOSAGE**

### **11 DESCRIPTION**

### **12 CLINICAL PHARMACOLOGY**

- 12.1 Mechanism of Action
- 12.3 Pharmacokinetics

### **13 NONCLINICAL TOXICOLOGY**

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

### **14 CLINICAL STUDIES**

- 14.1 Congestive Heart Failure Post-Myocardial Infarction
- 14.2 Hypertension

### **16 HOW SUPPLIED/STORAGE AND HANDLING**

### **17 PATIENT COUNSELING INFORMATION**

\* Sections or subsections omitted from the full prescribing information are not listed.

---

## **FULL PRESCRIBING INFORMATION**

### **1 INDICATIONS AND USAGE**

## 1.1 Patient Selection Considerations

Serum potassium levels should be measured before initiating eplerenone therapy, and eplerenone should not be prescribed if serum potassium is  $>5.5$  mEq/L. [See *CONTRAINDICATIONS (4)*].

## 1.2 Congestive Heart Failure Post-Myocardial Infarction

Eplerenone is indicated to improve survival of stable patients with left ventricular (LV) systolic dysfunction (ejection fraction  $\leq 40\%$ ) and clinical evidence of congestive heart failure (CHF) after an acute myocardial infarction (MI).

## 1.3 Hypertension

Eplerenone is indicated for the treatment of hypertension, to lower blood pressure. Lowering blood pressure reduces the risk of fatal and nonfatal cardiovascular (CV) events, primarily strokes and MI. These benefits have been seen in controlled trials of antihypertensive drugs from a wide variety of pharmacologic classes.

Control of high blood pressure should be part of comprehensive CV risk management, including, as appropriate, lipid control, diabetes management, antithrombotic therapy, smoking cessation, exercise, and limited sodium intake. Many patients will require more than one drug to achieve blood pressure goals. For specific advice on goals and management, see published guidelines, such as those of the National High Blood Pressure Education Program's Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure (JNC).

Numerous antihypertensive drugs, from a variety of pharmacologic classes and with different mechanisms of action, have been shown in randomized controlled trials to reduce CV morbidity and mortality, and it can be concluded that it is blood pressure reduction, and not some other pharmacologic property of the drugs, that is largely responsible for those benefits. The largest and most consistent CV outcome benefit has been a reduction in the risk of stroke, but reductions in MI and CV mortality also have been seen regularly.

Elevated systolic or diastolic pressure causes increased CV risk, and the absolute risk increase per mmHg is greater at higher blood pressures, so that even modest reductions of severe hypertension can provide substantial benefit. Relative risk reduction from blood pressure reduction is similar across populations with varying absolute risk, so the absolute benefit is greater in patients who are at higher risk independent of their hypertension (for example, patients with diabetes or hyperlipidemia), and such patients would be expected to benefit from more aggressive treatment to a lower blood pressure goal.

Some antihypertensive drugs have smaller blood pressure effects (as monotherapy) in black patients, and many antihypertensive drugs have additional approved indications and effects (e.g., on angina, heart failure, or diabetic kidney disease). These considerations may guide selection of therapy.

Eplerenone may be used alone or in combination with other antihypertensive agents.

## 2 DOSAGE AND ADMINISTRATION

### 2.1 Congestive Heart Failure Post-Myocardial Infarction

Treatment should be initiated at 25 mg once daily and titrated to the recommended dose of 50 mg once daily, preferably within 4 weeks as tolerated by the patient. Eplerenone may be administered with or without food.

Once treatment with eplerenone has begun, adjust the dose based on the serum potassium level as shown in Table 1.

**Table 1. Dose Adjustment in Congestive Heart Failure Post-MI**

<b>Serum</b>		
--------------	--	--

Potassium (mEq/L)	Action	Dose Adjustment
< 5.0	Increase	25 mg every other day to 25 mg once daily 25 mg once daily to 50 mg once daily
5.0–5.4	Maintain	No adjustment
5.5–5.9	Decrease	50 mg once daily to 25 mg once daily 25 mg once daily to 25 mg every other day 25 mg every other day to withhold
≥ 6.0	Withhold	Restart at 25 mg every other day when potassium levels fall to <5.5 mEq/L

## 2.2 Hypertension

The recommended starting dose of eplerenone is 50 mg administered once daily. The full therapeutic effect of eplerenone is apparent within 4 weeks. For patients with an inadequate blood pressure response to 50 mg once daily the dosage of eplerenone should be increased to 50 mg twice daily. Higher dosages of eplerenone are not recommended because they have no greater effect on blood pressure than 100 mg and are associated with an increased risk of hyperkalemia. [See *CLINICAL STUDIES* (14.2).]

## 2.3 Recommended Monitoring

Serum potassium should be measured before initiating eplerenone therapy, within the first week, and at one month after the start of treatment or dose adjustment. Serum potassium should be assessed periodically thereafter. Patient characteristics and serum potassium levels may indicate that additional monitoring is appropriate. [See *WARNINGS AND PRECAUTIONS* (5.1), *ADVERSE REACTIONS* (6.2).] In the EPHEsus study [See *CLINICAL STUDIES* (14.1)], the majority of hyperkalemia was observed within the first three months after randomization.

In all patients taking eplerenone who start taking a moderate CYP3A4 inhibitor, check serum potassium and serum creatinine in 3–7 days.

## 2.4 Dose Modifications for Specific Populations

For hypertensive patients receiving moderate CYP3A4 inhibitors (e.g., erythromycin, saquinavir, verapamil, and fluconazole), the starting dose of eplerenone should be reduced to 25 mg once daily. [See *DRUG INTERACTIONS* (7.1).]

No adjustment of the starting dose is recommended for the elderly or for patients with mild-to-moderate hepatic impairment. [See *CLINICAL PHARMACOLOGY* (12.3).]

## 3 DOSAGE FORMS AND STRENGTHS

- 25 mg tablets: yellow diamond biconvex film-coated tablets debossed with "G" on one side and "25mg" on the other
- 50 mg tablets: yellow diamond biconvex film-coated tablets debossed with "G" on one side and "50mg" on the other

## 4 CONTRAINDICATIONS

### For All Patients

Eplerenone is contraindicated in all patients with:

- serum potassium >5.5 mEq/L at initiation,
- creatinine clearance ≤30 mL/min, or
- concomitant administration of strong CYP3A4 inhibitors (e.g., ketoconazole, itraconazole,

nefazodone, troleandomycin, clarithromycin, ritonavir, and nelfinavir). [See *DRUG INTERACTIONS (7.1)*, *CLINICAL PHARMACOLOGY (12.3)*.]

#### For Patients Treated for Hypertension

Eplerenone is contraindicated for the treatment of hypertension in patients with:

- type 2 diabetes with microalbuminuria,
- serum creatinine >2.0 mg/dL in males or >1.8 mg/dL in females,
- creatinine clearance <50 mL/min, or
- concomitant administration of potassium supplements or potassium-sparing diuretics (e.g., amiloride, spironolactone, or triamterene). [See *WARNINGS AND PRECAUTIONS (5.1)*, *ADVERSE REACTIONS (6.2)*, *DRUG INTERACTIONS (7)*, and *CLINICAL PHARMACOLOGY (12.3)*.]

## **5 WARNINGS AND PRECAUTIONS**

### **5.1 Hyperkalemia**

Minimize the risk of hyperkalemia with proper patient selection and monitoring, and avoidance of certain concomitant medications [See *CONTRAINDICATIONS (4)*, *ADVERSE REACTIONS (6.2)*, and *DRUG INTERACTIONS (7)*]. Monitor patients for the development of hyperkalemia until the effect of eplerenone is established. Patients who develop hyperkalemia (>5.5 mEq/L) may continue eplerenone therapy with proper dose adjustment. Dose reduction decreases potassium levels. [See *DOSAGE AND ADMINISTRATION (2.1)*.]

The rates of hyperkalemia increase with declining renal function. [See *ADVERSE REACTIONS (6.2)*.] Patients with hypertension who have serum creatinine levels >2.0 mg/dL (males) or >1.8 mg/dL (females) or creatinine clearance ≤50 mL/min should not be treated with eplerenone. [See *CONTRAINDICATIONS (4)*.] Patients with CHF post-MI who have serum creatinine levels >2.0 mg/dL (males) or >1.8 mg/dL (females) or creatinine clearance ≤50 mL/min should be treated with eplerenone with caution.

Diabetic patients with CHF post-MI should also be treated with caution, especially those with proteinuria. The subset of patients in the EPHESUS study with both diabetes and proteinuria on the baseline urinalysis had increased rates of hyperkalemia compared to patients with either diabetes or proteinuria. [See *ADVERSE REACTIONS (6.2)*.]

The risk of hyperkalaemia may increase when eplerenone is used in combination with an angiotensin converting enzyme (ACE) inhibitor and/or an angiotensin receptor blocker (ARB). [See *DRUG INTERACTIONS (7.2)*.]

### **5.2 Impaired Hepatic Function**

Mild-to-moderate hepatic impairment did not increase the incidence of hyperkalemia. In 16 subjects with mild-to-moderate hepatic impairment who received 400 mg of eplerenone, no elevations of serum potassium above 5.5 mEq/L were observed. The mean increase in serum potassium was 0.12 mEq/L in patients with hepatic impairment and 0.13 mEq/L in normal controls. The use of eplerenone in patients with severe hepatic impairment has not been evaluated. [See *CLINICAL PHARMACOLOGY (12.3)*.]

### **5.3 Impaired Renal Function**

Patients with decreased renal function are at increased risk of hyperkalemia. [See *CONTRAINDICATIONS (4)*, *WARNINGS AND PRECAUTIONS (5.1)*, *ADVERSE REACTIONS (6.1)*.]

## **6 ADVERSE REACTIONS**

The following adverse reactions are discussed in greater detail in other sections of the labeling:

- Hyperkalemia [See WARNINGS AND PRECAUTIONS (5.1)]

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in clinical trials of another drug and may not reflect the rates observed in practice.

## 6.1 Clinical Trials Experience

### Congestive Heart Failure Post-Myocardial Infarction

In EPHESUS, safety was evaluated in 3307 patients treated with eplerenone and 3301 placebo-treated patients. The overall incidence of adverse events reported with eplerenone (78.9%) was similar to placebo (79.5%). Adverse events occurred at a similar rate regardless of age, gender, or race. Patients discontinued treatment due to an adverse event at similar rates in either treatment group (4.4% eplerenone vs. 4.3% placebo), with the most common reasons for discontinuation being hyperkalemia, MI, and abnormal renal function.

Adverse reactions that occurred more frequently in patients treated with eplerenone than placebo were hyperkalemia (3.4% vs. 2.0%) and increased creatinine (2.4% vs. 1.5%). Discontinuations due to hyperkalemia or abnormal renal function were less than 1.0% in both groups. Hypokalemia occurred less frequently in patients treated with eplerenone (0.6% vs. 1.6%).

The rates of sex hormone-related adverse events are shown in Table 2.

**Table 2. Rates of Sex Hormone-Related Adverse Events in EPHESUS**

	Rates in Males			Rates in Females
	Gynecomastia	Mastodynia	Either	Abnormal Vaginal Bleeding
Eplerenone	0.4%	0.1%	0.5%	0.4%
Placebo	0.5%	0.1%	0.6%	0.4%

### Hypertension

Eplerenone has been evaluated for safety in 3091 patients treated for hypertension. A total of 690 patients were treated for over 6 months and 106 patients were treated for over 1 year.

In placebo-controlled studies, the overall rates of adverse events were 47% with eplerenone and 45% with placebo. Adverse events occurred at a similar rate regardless of age, gender, or race. Therapy was discontinued due to an adverse event in 3% of patients treated with eplerenone and 3% of patients given placebo. The most common reasons for discontinuation of eplerenone were headache, dizziness, angina pectoris/MI, and increased GGT. The adverse events that were reported at a rate of at least 1% of patients and at a higher rate in patients treated with eplerenone in daily doses of 25 mg to 400 mg versus placebo are shown in Table 3.

**Table 3. Rates (%) of Adverse Events Occurring in Placebo-Controlled Hypertension Studies in ≥1% of Patients Treated with Eplerenone (25 mg to 400 mg) and at a More Frequent Rate than in Placebo-Treated Patients**

	Eplerenone (n=945)	Placebo (n=372)
<b>Metabolic</b>		
Hypercholesterolemia	1	0
Hypertriglyceridemia	1	0
<b>Digestive</b>		
Diarrhea	2	1

Abdominal pain	1	0
<b>Urinary</b>		
Albuminuria	1	0
<b>Respiratory</b>		
Coughing	2	1
<b>Central/Peripheral Nervous System</b>		
Dizziness	3	2
<b>Body as a Whole</b>		
Fatigue	2	1
Influenza-like symptoms	2	1

Note: Adverse events that are too general to be informative or are very common in the treated population are excluded.

Gynecomastia and abnormal vaginal bleeding were reported with eplerenone but not with placebo. The rates of these sex hormone-related adverse events are shown in Table 4. The rates increased slightly with increasing duration of therapy. In females, abnormal vaginal bleeding was also reported in 0.8% of patients on antihypertensive medications (other than spironolactone) in active control arms of the studies with eplerenone.

**Table 4. Rates of Sex Hormone-Related Adverse Events with Eplerenone in Hypertension Clinical Studies**

	Rates in Males			Rates in Females
	Gynecomastia	Mastodynia	Either	Abnormal Vaginal Bleeding
All controlled studies	0.5%	0.8%	1.0%	0.6%
Controlled studies lasting $\geq 6$ months	0.7%	1.3%	1.6%	0.8%
Open-label, long-term study	1.0%	0.3%	1.0%	2.1%

## 6.2 Clinical Laboratory Test Findings

### Congestive Heart Failure Post-Myocardial Infarction

**Creatinine:** Increases of more than 0.5 mg/dL were reported for 6.5% of patients administered eplerenone and for 4.9% of placebo-treated patients.

**Potassium:** In EPHESUS [see *CLINICAL STUDIES (14.1)*], the frequencies of patients with changes in potassium ( $<3.5$  mEq/L or  $>5.5$  mEq/L or  $\geq 6.0$  mEq/L) receiving eplerenone compared with placebo are displayed in Table 5.

**Table 5. Hypokalemia ( $<3.5$  mEq/L) or Hyperkalemia ( $>5.5$  or  $\geq 6.0$  mEq/L) in EPHESUS**

Potassium (mEq/L)	Eplerenone (N=3251) n (%)	Placebo (N=3237) n (%)
$< 3.5$	273 (8.4)	424 (13.1)
$>5.5$	508 (15.6)	363 (11.2)

≥ 6.0	180 (5.5)	126 (3.9)
-------	-----------	-----------

Table 6 shows the rates of hyperkalemia in EPHESUS as assessed by baseline renal function (creatinine clearance).

**Table 6. Rates of Hyperkalemia ( >5.5 mEq/L) in EPHESUS by Baseline Creatinine Clearance\***

<b>Baseline Creatinine Clearance</b>	<b>Eplerenone (N=508) n (%)</b>	<b>Placebo (N=363) n (%)</b>
≤30 mL/min	160 (32)	82 (23)
31–50 mL/min	122 (24)	46 (13)
51–70 mL/min	86 (17)	48 (13)
>70 mL/min	56 (11)	32 (9)

\* Estimated using the Cockcroft-Gault formula.

Table 7 shows the rates of hyperkalemia in EPHESUS as assessed by two baseline characteristics: presence/absence of proteinuria from baseline urinalysis and presence/absence of diabetes. [See *WARNINGS AND PRECAUTIONS (5.1).*]

**Table 7. Rates of Hyperkalemia ( >5.5 mEq/L) in EPHESUS by Proteinuria and History of Diabetes\***

	<b>Eplerenone (N=508) n (%)</b>	<b>Placebo (N=363) n (%)</b>
Proteinuria, no Diabetes	81 (16)	40 (11)
Diabetes, no Proteinuria	91 (18)	47 (13)
Proteinuria and Diabetes	132 (26)	58 (16)

\* Diabetes assessed as positive medical history at baseline; proteinuria assessed by positive dipstick urinalysis at baseline.

### Hypertension

**Potassium:** In placebo-controlled fixed-dose studies, the mean increases in serum potassium were dose-related and are shown in Table 8 along with the frequencies of values >5.5 mEq/L.

**Table 8. Increases in Serum Potassium in the Placebo-Controlled, Fixed-Dose Hypertension Studies of Eplerenone**

		<b>Mean Increase mEq/L</b>	<b>% &gt;5.5 mEq/L</b>
<b>Daily Dosage</b>	<b>n</b>		
Placebo	194	0	1
25	97	0.08	0
50	245	0.14	0
100	193	0.09	1
200	139	0.19	1
400	104	0.36	8.7

Patients with both type 2 diabetes and microalbuminuria are at increased risk of developing persistent



hyperkalemia. In a study of such patients taking eplerenone 200 mg, the frequencies of maximum serum potassium levels >5.5 mEq/L were 33% with eplerenone given alone and 38% when eplerenone was given with enalapril.

Rates of hyperkalemia increased with decreasing renal function. In all studies, serum potassium elevations >5.5 mEq/L were observed in 10.4% of patients treated with eplerenone with baseline calculated creatinine clearance <70 mL/min, 5.6% of patients with baseline creatinine clearance of 70 to 100 mL/min, and 2.6% of patients with baseline creatinine clearance of >100 mL/min. [See **WARNINGS AND PRECAUTIONS** (5.1).]

**Sodium:** Serum sodium decreased in a dose-related manner. Mean decreases ranged from 0.7 mEq/L at 50 mg daily to 1.7 mEq/L at 400 mg daily. Decreases in sodium (<135 mEq/L) were reported for 2.3% of patients administered eplerenone and 0.6% of placebo-treated patients.

**Triglycerides:** Serum triglycerides increased in a dose-related manner. Mean increases ranged from 7.1 mg/dL at 50 mg daily to 26.6 mg/dL at 400 mg daily. Increases in triglycerides (above 252 mg/dL) were reported for 15% of patients administered eplerenone and 12% of placebo-treated patients.

**Cholesterol:** Serum cholesterol increased in a dose-related manner. Mean changes ranged from a decrease of 0.4 mg/dL at 50 mg daily to an increase of 11.6 mg/dL at 400 mg daily. Increases in serum cholesterol values greater than 200 mg/dL were reported for 0.3% of patients administered eplerenone and 0% of placebo-treated patients.

**Liver Function Tests:** Serum alanine aminotransferase (ALT) and gamma glutamyl transpeptidase (GGT) increased in a dose-related manner. Mean increases ranged from 0.8 U/L at 50 mg daily to 4.8 U/L at 400 mg daily for ALT and 3.1 U/L at 50 mg daily to 11.3 U/L at 400 mg daily for GGT. Increases in ALT levels greater than 120 U/L (3 times upper limit of normal) were reported for 15/2259 patients administered eplerenone and 1/351 placebo-treated patients. Increases in ALT levels greater than 200 U/L (5 times upper limit of normal) were reported for 5/2259 of patients administered eplerenone and 1/351 placebo-treated patients. Increases of ALT greater than 120 U/L and bilirubin greater than 1.2 mg/dL were reported 1/2259 patients administered eplerenone and 0/351 placebo-treated patients. Hepatic failure was not reported in patients receiving eplerenone.

**BUN/Creatinine:** Serum creatinine increased in a dose-related manner. Mean increases ranged from 0.01 mg/dL at 50 mg daily to 0.03 mg/dL at 400 mg daily. Increases in blood urea nitrogen to greater than 30 mg/dL and serum creatinine to greater than 2 mg/dL were reported for 0.5% and 0.2%, respectively, of patients administered eplerenone and 0% of placebo-treated patients.

**Uric Acid:** Increases in uric acid to greater than 9 mg/dL were reported in 0.3% of patients administered eplerenone and 0% of placebo-treated patients.

### 6.3 Postmarketing Experience

The following adverse reactions have been identified during postapproval use of eplerenone. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

*Skin:* angioneurotic edema, rash

## 7 DRUG INTERACTIONS

### 7.1 CYP3A4 Inhibitors

Because eplerenone metabolism is predominantly mediated via CYP3A4, do not use eplerenone with drugs that are strong inhibitors of CYP3A4. [See **CONTRAINDICATIONS** (4) and **CLINICAL PHARMACOLOGY** (12.3).]

In patients with hypertension taking moderate CYP3A4 inhibitors, reduce the starting dose of eplerenone to 25 mg once daily. [See **DOSAGE AND ADMINISTRATION** (2.3, 2.4) and **CLINICAL**

## 7.2 ACE Inhibitors and Angiotensin II Receptor Antagonists

The risk of hyperkalaemia may increase when eplerenone is used in combination with an ACE inhibitor and/or an ARB. A close monitoring of serum potassium and renal function is recommended, especially in patients at risk for impaired renal function, e.g., the elderly. [See WARNINGS AND PRECAUTIONS (5.1)]

### Congestive Heart Failure Post-Myocardial Infarction

In EPHESUS [see CLINICAL STUDIES (14.1)], 3020 (91%) patients receiving eplerenone 25 mg to 50 mg also received ACE inhibitors or angiotensin II receptor antagonists (ACEI/ARB). Rates of patients with maximum potassium levels >5.5 mEq/L were similar regardless of the use of ACEI/ARB.

### Hypertension

In clinical studies of patients with hypertension, the addition of eplerenone 50 mg to 100 mg to ACE inhibitors and ARB increased mean serum potassium slightly (about 0.09–0.13 mEq/L). In a study in diabetics with microalbuminuria, eplerenone 200 mg combined with the ACE inhibitor enalapril 10 mg increased the frequency of hyperkalemia (serum potassium >5.5 mEq/L) from 17% on enalapril alone to 38%.

## 7.3 Lithium

A drug interaction study of eplerenone with lithium has not been conducted. Lithium toxicity has been reported in patients receiving lithium concomitantly with diuretics and ACE inhibitors. Serum lithium levels should be monitored frequently if eplerenone is administered concomitantly with lithium.

## 7.4 Nonsteroidal Anti-Inflammatory Drugs (NSAIDs)

A drug interaction study of eplerenone with an NSAID has not been conducted. The administration of other potassium-sparing antihypertensives with NSAIDs has been shown to reduce the antihypertensive effect in some patients and result in severe hyperkalemia in patients with impaired renal function. Therefore, when eplerenone and NSAIDs are used concomitantly, patients should be observed to determine whether the desired effect on blood pressure is obtained and monitored for changes in serum potassium levels.

## 8 USE IN SPECIFIC POPULATIONS

### 8.1 Pregnancy

Pregnancy Category B

There are no adequate and well-controlled studies in pregnant women. Eplerenone should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

### Teratogenic Effects

Embryo-fetal development studies were conducted with doses up to 1000 mg/kg/day in rats and 300 mg/kg/day in rabbits (exposures up to 32 and 31 times the human AUC for the 100 mg/day therapeutic dose, respectively). No teratogenic effects were seen in rats or rabbits, although decreased body weight in maternal rabbits and increased rabbit fetal resorptions and post-implantation loss were observed at the highest administered dosage. Because animal reproduction studies are not always predictive of human response, eplerenone should be used during pregnancy only if clearly needed.

### 8.3 Nursing Mothers

The concentration of eplerenone in human breast milk after oral administration is unknown. However, preclinical data show that eplerenone and/or metabolites are present in rat breast milk (0.85:1

[milk:plasma] AUC ratio) obtained after a single oral dose. Peak concentrations in plasma and milk were obtained from 0.5 to 1 hour after dosing. Rat pups exposed by this route developed normally. Because many drugs are excreted in human milk and because of the unknown potential for adverse effects on the nursing infant, a decision should be made whether to discontinue nursing or discontinue the drug, taking into account the importance of the drug to the mother.

#### **8.4 Pediatric Use**

In a 10-week study of 304 hypertensive pediatric patients age 4 to 17 years treated with eplerenone up to 100 mg per day, doses that produced exposure similar to that in adults, eplerenone did not lower blood pressure effectively. In this study and in a 1-year pediatric safety study in 149 patients, the incidence of reported adverse events was similar to that of adults.

Eplerenone has not been studied in hypertensive patients less than 4 years old because the study in older pediatric patients did not demonstrate effectiveness.

Eplerenone has not been studied in pediatric patients with heart failure.

#### **8.5 Geriatric Use**

##### *Congestive Heart Failure Post-Myocardial Infarction*

Of the total number of patients in EPHESUS, 3340 (50%) were 65 and over, while 1326 (20%) were 75 and over. Patients greater than 75 years did not appear to benefit from the use of eplerenone. [See *CLINICAL STUDIES (14.1).*]

No differences in overall incidence of adverse events were observed between elderly and younger patients. However, due to age-related decreases in creatinine clearance, the incidence of laboratory-documented hyperkalemia was increased in patients 65 and older. [See *WARNINGS AND PRECAUTIONS (5.1).*]

##### *Hypertension*

Of the total number of subjects in clinical hypertension studies of eplerenone, 1123 (23%) were 65 and over, while 212 (4%) were 75 and over. No overall differences in safety or effectiveness were observed between elderly subjects and younger subjects.

### **10 OVERDOSAGE**


No cases of human overdosage with eplerenone have been reported. Lethality was not observed in mice, rats, or dogs after single oral doses that provided  $C_{\max}$  exposures at least 25 times higher than in humans receiving eplerenone 100 mg/day. Dogs showed emesis, salivation, and tremors at a  $C_{\max}$  41 times the human therapeutic  $C_{\max}$ , progressing to sedation and convulsions at higher exposures.

The most likely manifestation of human overdosage would be anticipated to be hypotension or hyperkalemia. Eplerenone cannot be removed by hemodialysis. Eplerenone has been shown to bind extensively to charcoal. If symptomatic hypotension should occur, supportive treatment should be instituted. If hyperkalemia develops, standard treatment should be initiated.

### **11 DESCRIPTION**

Eplerenone tablets contain eplerenone, a blocker of aldosterone binding at the mineralocorticoid receptor.

Eplerenone is chemically described as Pregn-4-ene-7,21-dicarboxylic acid, 9,11-epoxy-17-hydroxy-3-oxo-,  $\gamma$ -lactone, methyl ester, (7 $\alpha$ ,11 $\alpha$ ,17 $\alpha$ )-. Its empirical formula is  $C_{24}H_{30}O_6$  and it has a molecular weight of 414.50. The structural formula of eplerenone is represented below:



**eplerenone**

Eplerenone is an odorless, white to off-white crystalline powder. It is very slightly soluble in water, with its solubility essentially pH-independent. The octanol/water partition coefficient of eplerenone is approximately 7.1 at pH 7.0.

Eplerenone tablets for oral administration contain 25 mg or 50 mg of eplerenone and the following inactive ingredients: lactose, microcrystalline cellulose, croscarmellose sodium, hypromellose, sodium lauryl sulfate, talc, magnesium stearate, titanium dioxide, polyethylene glycol, polysorbate 80, and iron oxide yellow and iron oxide red.

## **12 CLINICAL PHARMACOLOGY**

### **12.1 Mechanism of Action**

Eplerenone binds to the mineralocorticoid receptor and blocks the binding of aldosterone, a component of the renin-angiotensin-aldosterone-system (RAAS). Aldosterone synthesis, which occurs primarily in the adrenal gland, is modulated by multiple factors, including angiotensin II and non-RAAS mediators such as adrenocorticotrophic hormone (ACTH) and potassium. Aldosterone binds to mineralocorticoid receptors in both epithelial (e.g., kidney) and non-epithelial (e.g., heart, blood vessels, and brain) tissues and increases blood pressure through induction of sodium reabsorption and possibly other mechanisms.

Eplerenone has been shown to produce sustained increases in plasma renin and serum aldosterone, consistent with inhibition of the negative regulatory feedback of aldosterone on renin secretion. The resulting increased plasma renin activity and aldosterone circulating levels do not overcome the effects of eplerenone.

Eplerenone selectively binds to recombinant human mineralocorticoid receptors relative to its binding to recombinant human glucocorticoid, progesterone, and androgen receptors.

### **12.3 Pharmacokinetics**

Eplerenone is cleared predominantly by cytochrome P450 (CYP) 3A4 metabolism, with an elimination half-life of 4 to 6 hours. Steady state is reached within 2 days. Absorption is not affected by food. Inhibitors of CYP3A4 (e.g., ketoconazole, saquinavir) increase blood levels of eplerenone.

#### *Absorption and Distribution*

Mean peak plasma concentrations of eplerenone are reached approximately 1.5 hours following oral administration. The absolute bioavailability of eplerenone is 69% following administration of a 100 mg

oral tablet. Both peak plasma levels ( $C_{\max}$ ) and area under the curve (AUC) are dose proportional for doses of 25 mg to 100 mg and less than proportional at doses above 100 mg.

The plasma protein binding of eplerenone is about 50% and it is primarily bound to alpha 1-acid glycoproteins. The apparent volume of distribution at steady state ranged from 43 to 90 L. Eplerenone does not preferentially bind to red blood cells.

#### Metabolism and Excretion

Eplerenone metabolism is primarily mediated via CYP3A4. No active metabolites of eplerenone have been identified in human plasma.

Less than 5% of an eplerenone dose is recovered as unchanged drug in the urine and feces. Following a single oral dose of radiolabeled drug, approximately 32% of the dose was excreted in the feces and approximately 67% was excreted in the urine. The elimination half-life of eplerenone is approximately 4 to 6 hours. The apparent plasma clearance is approximately 10 L/hr.

#### Age, Gender, and Race

The pharmacokinetics of eplerenone at a dose of 100 mg once daily has been investigated in the elderly ( $\geq 65$  years), in males and females, and in Blacks. At steady state, elderly subjects had increases in  $C_{\max}$  (22%) and AUC (45%) compared with younger subjects (18 to 45 years). The pharmacokinetics of eplerenone did not differ significantly between males and females. At steady state,  $C_{\max}$  was 19% lower and AUC was 26% lower in Blacks. [See *DOSAGE AND ADMINISTRATION* (2.4) and *USE IN SPECIFIC POPULATIONS* (8.5).]

#### Renal Impairment

The pharmacokinetics of eplerenone was evaluated in patients with varying degrees of renal impairment and in patients undergoing hemodialysis. Compared with control subjects, steady state AUC and  $C_{\max}$  were increased by 38% and 24%, respectively, in patients with severe renal impairment and were decreased by 26% and 3%, respectively, in patients undergoing hemodialysis. No correlation was observed between plasma clearance of eplerenone and creatinine clearance. Eplerenone is not removed by hemodialysis. [See *WARNINGS AND PRECAUTIONS* (5.3).]

#### Hepatic Impairment

The pharmacokinetics of eplerenone 400 mg has been investigated in patients with moderate (Child-Pugh Class B) hepatic impairment and compared with normal subjects. Steady state  $C_{\max}$  and AUC of eplerenone were increased by 3.6% and 42%, respectively.

#### Heart Failure

The pharmacokinetics of eplerenone 50 mg was evaluated in 8 patients with heart failure (NYHA classification II–IV) and 8 matched (gender, age, weight) healthy controls. Compared with the controls, steady state AUC and  $C_{\max}$  in patients with stable heart failure were 38% and 30% higher, respectively.

#### Drug-Drug Interactions [See *DRUG INTERACTIONS* (7).]

Eplerenone is metabolized primarily by CYP3A4. Inhibitors of CYP3A4 cause increased exposure [see *DRUG INTERACTIONS* (7.1)].

Drug-drug interaction studies were conducted with a 100 mg dose of eplerenone.

A pharmacokinetic study evaluating the administration of a single dose of eplerenone 100 mg with ketoconazole 200 mg two times a day, a strong inhibitor of the CYP3A4 pathway, showed a 1.7-fold increase in  $C_{\max}$  of eplerenone and a 5.4-fold increase in AUC of eplerenone.

Administration of eplerenone with moderate CYP3A4 inhibitors (e.g., erythromycin 500 mg BID, verapamil 240 mg once daily, saquinavir 1200 mg three times a day, fluconazole 200 mg once daily) resulted in increases in  $C_{\max}$  of eplerenone ranging from 1.4- to 1.6-fold and AUC from 2.0- to 2.9-fold.

Grapefruit juice caused only a small increase (about 25%) in exposure.

Eplerenone is not an inhibitor of CYP1A2, CYP3A4, CYP2C19, CYP2C9, or CYP2D6. Eplerenone did not inhibit the metabolism of amiodarone, amlodipine, astemizole, chlorzoxazone, cisapride, dexamethasone, dextromethorphan, diclofenac, 17 $\alpha$ -ethinyl estradiol, fluoxetine, losartan, lovastatin, mephobarbital, methylphenidate, methylprednisolone, metoprolol, midazolam, nifedipine, phenacetin, phenytoin, simvastatin, tolbutamide, triazolam, verapamil, or warfarin in vitro. Eplerenone is not a substrate or an inhibitor of P-Glycoprotein at clinically relevant doses.

No clinically significant drug-drug pharmacokinetic interactions were observed when eplerenone was administered with cisapride, cyclosporine, digoxin, glyburide, midazolam, oral contraceptives (norethindrone/ethinyl estradiol), simvastatin, or warfarin. St. John's wort (a CYP3A4 inducer) caused a small (about 30%) decrease in eplerenone AUC.

No significant changes in eplerenone pharmacokinetics were observed when eplerenone was administered with aluminum- and magnesium-containing antacids.

## **13 NONCLINICAL TOXICOLOGY**

### **13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility**

Eplerenone was non-genotoxic in a battery of assays including in vitro bacterial mutagenesis (Ames test in *Salmonella* spp. and *E. Coli*), in vitro mammalian cell mutagenesis (mouse lymphoma cells), in vitro chromosomal aberration (Chinese hamster ovary cells), in vivo rat bone marrow micronucleus formation, and in vivo/ex vivo unscheduled DNA synthesis in rat liver.

There was no drug-related tumor response in heterozygous P53 deficient mice when tested for 6 months at dosages up to 1000 mg/kg/day (systemic AUC exposures up to 9 times the exposure in humans receiving the 100 mg/day therapeutic dose). Statistically significant increases in benign thyroid tumors were observed after 2 years in both male and female rats when administered eplerenone 250 mg/kg/day (highest dose tested) and in male rats only at 75 mg/kg/day. These dosages provided systemic AUC exposures approximately 2 to 12 times higher than the average human therapeutic exposure at 100 mg/day. Repeat dose administration of eplerenone to rats increases the hepatic conjugation and clearance of thyroxine, which results in increased levels of TSH by a compensatory mechanism. Drugs that have produced thyroid tumors by this rodent-specific mechanism have not shown a similar effect in humans.

Male rats treated with eplerenone at 1000 mg/kg/day for 10 weeks (AUC 17 times that at the 100 mg/day human therapeutic dose) had decreased weights of seminal vesicles and epididymides and slightly decreased fertility. Dogs administered eplerenone at dosages of 15 mg/kg/day and higher (AUC 5 times that at the 100 mg/day human therapeutic dose) had dose-related prostate atrophy. The prostate atrophy was reversible after daily treatment for 1 year at 100 mg/kg/day. Dogs with prostate atrophy showed no decline in libido, sexual performance, or semen quality. Testicular weight and histology were not affected by eplerenone in any test animal species at any dosage.

## **14 CLINICAL STUDIES**

### **14.1 Congestive Heart Failure Post-Myocardial Infarction**

The eplerenone post-acute myocardial infarction heart failure efficacy and survival study (EPHESUS) was a multinational, multicenter, double-blind, randomized, placebo-controlled study in patients clinically stable 3 to 14 days after an acute MI with LV dysfunction (as measured by left ventricular ejection fraction [LVEF]  $\leq 40\%$ ) and either diabetes or clinical evidence of CHF (pulmonary congestion by exam or chest x-ray or S<sub>3</sub>). Patients with CHF of valvular or congenital etiology, patients with unstable post-infarct angina, and patients with serum potassium  $>5.0$  mEq/L or serum creatinine  $>2.5$  mg/dL were to be excluded. Patients were allowed to receive standard post-MI drug therapy and to

undergo revascularization by angioplasty or coronary artery bypass graft surgery.

Patients randomized to eplerenone were given an initial dose of 25 mg once daily and titrated to the target dose of 50 mg once daily after 4 weeks if serum potassium was < 5.0 mEq/L. Dosage was reduced or suspended anytime during the study if serum potassium levels were  $\geq 5.5$  mEq/L. [See *DOSAGE AND ADMINISTRATION (2.1).*]

EPHESUS randomized 6,632 patients (9.3% U.S.) at 671 centers in 27 countries. The study population was primarily white (90%, with 1% Black, 1% Asian, 6% Hispanic, 2% other) and male (71%). The mean age was 64 years (range, 22 to 94 years). The majority of patients had pulmonary congestion (75%) by exam or x-ray and were Killip Class II (64%). The mean ejection fraction was 33%. The average time to enrollment was 7 days post-MI. Medical histories prior to the index MI included hypertension (60%), coronary artery disease (62%), dyslipidemia (48%), angina (41%), type 2 diabetes (30%), previous MI (27%), and CHF (15%).


The mean dose of eplerenone was 43 mg/day. Patients also received standard care including aspirin (92%), ACE inhibitors (90%), beta-blockers (83%), nitrates (72%), loop diuretics (66%), or HMG-CoA reductase inhibitors (60%).

Patients were followed for an average of 16 months (range, 0 to 33 months). The ascertainment rate for vital status was 99.7%.

The co-primary endpoints for EPHESUS were (1) the time to death from any cause, and (2) the time to first occurrence of either cardiovascular mortality [defined as sudden cardiac death or death due to progression of CHF, stroke, or other CV causes] or CV hospitalization (defined as hospitalization for progression of CHF, ventricular arrhythmias, acute MI, or stroke).

For the co-primary endpoint for death from any cause, there were 478 deaths in the eplerenone group (14.4%) and 554 deaths in the placebo group (16.7%). The risk of death with eplerenone was reduced by 15% [hazard ratio equal to 0.85 (95% confidence interval 0.75 to 0.96;  $p = 0.008$  by log rank test)]. Kaplan-Meier estimates of all-cause mortality are shown in Figure 1 and the components of mortality are provided in Table 9.

Figure 1. Kaplan-Meier Estimates of All-Cause Mortality



**Table 9. Components of All-Cause Mortality in EPHESUS**

	<b>Eplerenone (N=3319) n (%)</b>	<b>Placebo (N=3313) n (%)</b>	<b>Hazard Ratio</b>	<b>p-value</b>
Death from any cause	478 (14.4)	554 (16.7)	0.85	0.008
CV Death	407 (12.3)	483 (14.6)	0.83	0.005
Non-CV Death	60 (1.8)	54 (1.6)		
Unknown or unwitnessed death	11 (0.3)	17 (0.5)		

Most CV deaths were attributed to sudden death, acute MI, and CHF.

The time to first event for the co-primary endpoint of CV death or hospitalization, as defined above, was longer in the eplerenone group (hazard ratio 0.87, 95% confidence interval 0.79 to 0.95,  $p = 0.002$ ). An analysis that included the time to first occurrence of CV mortality and all CV hospitalizations (atrial arrhythmia, angina, CV procedures, progression of CHF, MI, stroke, ventricular arrhythmia, or other CV causes) showed a smaller effect with a hazard ratio of 0.92 (95% confidence interval 0.86 to 0.99;  $p = 0.028$ ). The combined endpoints, including combined all-cause hospitalization and mortality were driven primarily by CV mortality. The combined endpoints in EPHESUS, including all-cause hospitalization and all-cause mortality, are presented in Table 10.

**Table 10. Rates of Death or Hospitalization in EPHESUS**



<b>Event</b>	<b>Eplerenone n (%)</b>	<b>Placebo n (%)</b>
CV death or hospitalization for progression of CHF, stroke, MI or ventricular arrhythmia*	885 (26.7)	993 (30.0)
Death	407 (12.3)	483 (14.6)
Hospitalization	606 (18.3)	649 (19.6)
CV death or hospitalization for progression of CHF, stroke, MI, ventricular arrhythmia, atrial arrhythmia, angina, CV procedures, or other CV causes (PVD; Hypotension)	1516 (45.7)	1610 (48.6)
Death	407 (12.3)	483 (14.6)
Hospitalization	1281 (38.6)	1307 (39.5)
All-cause death or hospitalization	1734 (52.2)	1833 (55.3)
Death*	478 (14.4)	554 (16.7)
Hospitalization	1497 (45.1)	1530 (46.2)


\* Co-Primary Endpoint.

Mortality hazard ratios varied for some subgroups as shown in Figure 2. Mortality hazard ratios appeared favorable for eplerenone for both genders and for all races or ethnic groups, although the numbers of non-Caucasians were low (648, 10%). Patients with diabetes without clinical evidence of CHF and patients greater than 75 years did not appear to benefit from the use of eplerenone. Such subgroup analyses must be interpreted cautiously.

**Figure 2. Hazard Ratios of All-Cause Mortality by Subgroups**








Analyses conducted for a variety of CV biomarkers did not confirm a mechanism of action by which mortality was reduced.

## 14.2 Hypertension


The safety and efficacy of eplerenone have been evaluated alone and in combination with other antihypertensive agents in clinical studies of 3091 hypertensive patients. The studies included 46% women, 14% Blacks, and 22% elderly (age  $\geq 65$ ). The studies excluded patients with elevated baseline serum potassium ( $>5.0$  mEq/L) and elevated baseline serum creatinine (generally  $>1.5$  mg/dL in males and  $>1.3$  mg/dL in females).

Two fixed-dose, placebo-controlled, 8- to 12-week monotherapy studies in patients with baseline diastolic blood pressures of 95 to 114 mm Hg were conducted to assess the antihypertensive effect of eplerenone. In these two studies, 611 patients were randomized to eplerenone and 140 patients to placebo. Patients received eplerenone in doses of 25 mg to 400 mg daily as either a single daily dose or divided into two daily doses. The mean placebo-subtracted reductions in trough cuff blood pressure achieved by eplerenone in these studies at doses up to 200 mg are shown in Figures 3 and 4.

**Figure 3. Eplerenone Dose Response - Trough Cuff SBP**  
Placebo-Subtracted Adjusted Mean Change from Baseline  
in Hypertension Studies



**Figure 4. Eplerenone Dose Response - Trough Cuff DBP**  
Placebo-Subtracted Adjusted Mean Change from Baseline  
in Hypertension Studies



Patients treated with eplerenone 50 mg to 200 mg daily experienced significant decreases in sitting systolic and diastolic blood pressure at trough with differences from placebo of 6–13 mm Hg (systolic) and 3–7 mm Hg (diastolic). These effects were confirmed by assessments with 24-hour ambulatory blood pressure monitoring (ABPM). In these studies, assessments of 24-hour ABPM data demonstrated that eplerenone, administered once or twice daily, maintained antihypertensive efficacy over the entire dosing interval. However, at a total daily dose of 100 mg, eplerenone administered as 50 mg twice per day produced greater trough cuff (4/3 mm Hg) and ABPM (2/1 mm Hg) blood pressure reductions than 100 mg given once daily.

Blood pressure lowering was apparent within 2 weeks from the start of therapy with eplerenone, with maximal antihypertensive effects achieved within 4 weeks. Stopping eplerenone following treatment for 8 to 24 weeks in six studies did not lead to adverse event rates in the week following withdrawal of eplerenone greater than following placebo or active control withdrawal. Blood pressures in patients not taking other antihypertensives rose 1 week after withdrawal of eplerenone by about 6/3 mm Hg, suggesting that the antihypertensive effect of eplerenone was maintained through 8 to 24 weeks.

Blood pressure reductions with eplerenone in the two fixed-dose monotherapy studies and other studies using titrated doses, as well as concomitant treatments, were not significantly different when analyzed by age, gender, or race with one exception. In a study in patients with low renin hypertension, blood pressure reductions in Blacks were smaller than those in whites during the initial titration period with eplerenone.

Eplerenone has been studied concomitantly with treatment with ACE inhibitors, ARB, calcium channel blockers, beta-blockers, and hydrochlorothiazide. When administered concomitantly with one of these drugs eplerenone usually produced its expected antihypertensive effects.

There was no significant change in average heart rate among patients treated with eplerenone in the combined clinical studies. No consistent effects of eplerenone on heart rate, QRS duration, or PR or QT interval were observed in 147 normal subjects evaluated for electrocardiographic changes during pharmacokinetic studies.

## 16 HOW SUPPLIED/STORAGE AND HANDLING

Eplerenone tablets, 25 mg, are yellow diamond biconvex film-coated tablets. They are debossed with "G" on one side and "25mg" on the other. They are supplied as follows:

NDC Number	Size
59762-1710-2	Bottle of 30 tablets
59762-1710-3	Bottle of 90 tablets
59762-1710-1	Hospital Unit Dose (10×10)

Eplerenone tablets, 50 mg, are yellow diamond biconvex film-coated tablets. They are debossed with "G" on one side and "50mg" on the other. They are supplied as follows:

NDC Number	Size
59762-1720-1	Bottle of 30 tablets
59762-1720-2	Bottle of 90 tablets

Store at 25°C (77°F); excursions permitted to 15–30°C (59–86°F) [See USP Controlled Room Temperature].

## 17 PATIENT COUNSELING INFORMATION

Patients receiving eplerenone should be informed:

- Not to use strong CYP3A4 inhibitors, such as ketoconazole, clarithromycin, nefazodone, ritonavir, and nelfinavir.
- Not to use potassium supplements or salt substitutes containing potassium without consulting the prescribing physician.
- To call their physician if they experience dizziness, diarrhea, vomiting, rapid or irregular heartbeat, lower extremity edema, or difficulty breathing.
- That periodic monitoring of blood pressure and serum potassium is important.

[See CONTRAINDICATIONS (4); WARNINGS AND PRECAUTIONS (5.1); and DRUG INTERACTIONS (7).]

LAB-0376-6.0

**PRINCIPAL DISPLAY PANEL - 25 mg Tablet Bottle Label**

**NDC 59762-1710-2**


30 Tablets

**GREENSTONE® BRAND**

**eplerenone  
tablets**

**25 mg**

**Rx only**




**PRINCIPAL DISPLAY PANEL - 25 mg Tablet Blister Pack**

13884600

**eplerenone  
tablet**

**25 mg**

Greenstone LLC  
Peapack, NJ 07977



## PRINCIPAL DISPLAY PANEL - 25 mg Tablet Blister Pack Carton

UNIT DOSE

NDC 59762-1710-1

100 Tablets


GREENSTONE® BRAND

**epplerenone**  
**tablets**

**25 mg**

For in-institution use only

**Rx only**



## PRINCIPAL DISPLAY PANEL - 50 mg Tablet Bottle Label

NDC 59762-1720-1

30 Tablets

GREENSTONE® BRAND


**epplerenone**  
**tablets**

50 mg  
Rx only

Store at 25°C (77°F); excursions permitted to 15-30°C (59-86°F) [see USP Controlled Room Temperature].

**DOSAGE AND USE**  
See accompanying prescribing information.

Each tablet contains 50 mg eplerenone.

  
Distributed by:  
**Greenstone LLC**  
Peapack, NJ 07977

NDC 59762-1720-1  
30 Tablets

GREENSTONE® BRAND

**eplerenone**  
tablets

**50 mg**

Rx only

UPC/EAN  
FPO (80% x 5.5mm)  
N 3 59762-1720-1 3  
13885200

EPLERENONE

eplerenone tablet, film coated

Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:59 762-1710
Route of Administration	ORAL	DEA Schedule	
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
EPLERENONE (UNII: 6995V82D0B) (EPLERENONE - UNII:6995V82D0B)		EPLERENONE	25 mg
Inactive Ingredients			
Ingredient Name			Strength
LACTOSE (UNII: J2B2A4N98G)			
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)			
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)			
HYPROMELLOSES (UNII: 3NXW29V3WO)			
SODIUM LAURYL SULFATE (UNII: 368GB5141J)			
TALC (UNII: 7SEV7J4R1U)			
MAGNESIUM STEARATE (UNII: 70097M6I3O)			
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)			
POLYETHYLENE GLYCOLS (UNII: 3WJQ0SDW1A)			
POLYSORBATE 80 (UNII: 6OZP39ZG8H)			



FERRIC OXIDE YELLOW (UNII: EX438O2MRT)				
FERRIC OXIDE RED (UNII: 1K09F3G675)				
Product Characteristics				
Color	YELLOW	Score	no score	
Shape	DIAMOND (biconvex)	Size	7mm	
Flavor		Imprint Code	G;25mg	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:59762-1710-2	30 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:59762-1710-3	90 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:59762-1710-1	100 in 1 CARTON		
3		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA AUTHORIZED GENERIC		NDA021437	09/27/2002	

## EPLERENONE

eplerenone tablet, film coated

<b>Product Information</b>			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:59762-1720
Route of Administration	ORAL	DEA Schedule	
<b>Active Ingredient/Active Moiety</b>			
Ingredient Name		Basis of Strength	Strength
EPLERENONE (UNII: 6995V82D0B) (EPLERENONE - UNII:6995V82D0B)		EPLERENONE	50 mg
<b>Inactive Ingredients</b>			
Ingredient Name			Strength
LACTOSE (UNII: J2B2A4N98G)			
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)			
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)			
HYPROMELLOSES (UNII: 3NXW29V3WO)			
SODIUM LAURYL SULFATE (UNII: 368GB5141J)			
TALC (UNII: 7SEV7J4R1U)			
MAGNESIUM STEARATE (UNII: 70097M6I3O)			
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)			

POLYETHYLENE GLYCOLS (UNII: 3WJQ0SDW1A)				
POLYSORBATE 80 (UNII: 6OZP39ZG8H)				
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)				
FERRIC OXIDE RED (UNII: 1K09F3G675)				
Product Characteristics				
Color	YELLOW	Score	no score	
Shape	DIAMOND (biconvex)	Size	9mm	
Flavor		Imprint Code	G;50mg	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:59762-1720-1	30 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:59762-1720-2	90 in 1 BOTTLE; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA AUTHORIZED GENERIC		NDA021437	09/27/2002	

**Labeler** - Greenstone LLC (825560733)

## Establishment

Name	Address	ID/FEI	Business Operations
Pharmacia and Upjohn Company		829076566	ANALYSIS(59762-1720, 59762-1710) , MANUFACTURE(59762-1720, 59762-1710) , PACK(59762-1720, 59762-1710)

## Establishment

Name	Address	ID/FEI	Business Operations
Pfizer Pharmaceuticals LLC		829084552	ANALYSIS(59762-1720, 59762-1710) , LABEL(59762-1720, 59762-1710) , PACK(59762-1720, 59762-1710)

## Establishment

Name	Address	ID/FEI	Business Operations
Excella GmbH		329809800	API MANUFACTURE(59762-1720, 59762-1710)